

Prescription Galactagogues

A galactagogue is an herb or medication taken by a breastfeeding mother to increase her milk supply. Throughout history and in most countries, women have turned to plants or herbs to increase their milk supply. (*See our information on **Herbal Galactagogues.***) There are two **prescription medications** that are known to increase serum prolactin levels (the hormone connected to making milk): **Reglan and Domperidone.**

Not all women can increase their low milk supply with a galactagogue due to the many underlying causes of low milk production. Prior to taking a galactagogue, MilkWorks highly recommends a **feeding assessment** by an IBCLC or breastfeeding medicine specialist to identify possible causes and solutions.

Unfortunately, there is no magic pill when it comes to increasing milk supply. **Frequent milk removal is still the most effective way to increase your milk.**

REGLAN (metoclopramide) is a stomach medication that may increase milk supply. However, it commonly causes depression and fatigue, and may cause involuntary muscle spasms, called tardive dyskinesia. **As a result, we rarely recommend using it.**

- There is no official established dosage for prescribing Reglan to increase milk supply. Most studies have administered 5 - 10 mg per dose 2 or 3 times a day for 7 to 14 days. Some studies have used a tapering dose for the last few days of treatment, to avoid an abrupt drop in milk supply after discontinuing the drug. No published literature supports the efficacy or safety of higher doses, longer treatment periods, or repeated courses of therapy.

DOMPERIDONE is also a stomach medication. It is not approved for marketing in the United States by the Food and Drug Administration because it is not manufactured in the U.S. It is available in Canada and other countries. Nebraska's Attorney General investigated practitioners who were prescribing domperidone via compounding pharmacies in 2006 and claimed that **prescribing this medication is illegal as it is not an FDA approved drug.** The FDA also released a statement of concern about cardiac arrhythmias. If mothers attempt to purchase this medication from outside the country, the package may be confiscated at the border.

- The typical dosage for domperidone is 10 mg 3 times a day. Two small studies found no statistically significant additional increase in milk output with a dosage of 20 mg 3 times a day. Women who failed to respond to the lower dose did not respond to the higher dose either. Doses >30 mg a day may increase the risk of arrhythmia and sudden cardiac death in patients receiving domperidone.

For additional information about galactagogues:

- Please read the Academy of Breastfeeding Medicine's protocol at <https://www.bfmed.org/> OR
- Visit LACTMED, a free online National of Library of Medicine data base at <https://toxnet.nlm.nih.gov/>

Reviewed: copyright January 2021

MilkWorks



Community
Breastfeeding
Center

5930 S. 58th Street
(in the Trade Center)
Lincoln, NE 68516
(402) 423-6402

10818 Elm Street
Rockbrook Village
Omaha, NE 68144
(402) 502-0617

For additional
information:
www.milkworks.org