

Anesthesia & Pain Medications for Breastfeeding Mothers

How soon is it safe to breastfeed if you receive a local or general anesthesia, or if you need to take a pain relief medication?

Anesthesia

- **General Anesthesia** is designed to wear off quickly. As soon as you are awake and alert, it has cleared from your system enough to breastfeed, assuming your baby is healthy and full term (at last 39 weeks gestation). If your baby is premature or ill, ask your baby's doctor how long you should wait before breastfeeding. If your baby is prone to apnea, low blood pressure, or low muscle tone, it may be best to pump and discard your milk for 6 – 12 hours before you resume breastfeeding.
- **If you receive a single injection for a dental procedure for sedation or pain**, you can breastfeed as soon as you are alert and stable. Although short-acting agents such as fentanyl and midazolam are preferred, single doses of meperidine/pethidine or diazepam are unlikely to affect a breastfeeding baby.
- **If you are having plastic surgery**, such as liposuction, where large doses of local anesthetics (lidocaine/xylocaine or lignocaine) are used, you should probably pump and discard your milk for 12 hours before you resume breastfeeding.
- **Drugs used for anesthetic induction**, such as propofol, midazolam, etomidate, or thiopental, enter the milk minimally because they are cleared from the blood so quickly.
- There is little research on **anesthetic gases**, but they are only in the blood briefly and are unlikely to be in breastmilk in significant amounts.

Pain Medications

- **Morphine** is considered an acceptable pain medicine for breastfeeding mothers because it does not get into breastmilk well and is not absorbed well by a baby.
- **Meperidine (Demerol) should NOT be given to breastfeeding mothers**, as it is more likely to cause sleepiness and depressed breathing in a baby.
- **Fentanyl** levels in breast milk have been studied and are extremely low after two hours.
- **Remifentanyl (Ultiva)** is a VERY short acting drug. Milk levels have not been studied, but it should not cause trouble for a baby.
- **Hydrocodone** rarely causes excessive sleepiness in babies. Take the smallest effective dose and stop taking it as soon as you can control your pain with Tylenol or Ibuprofen.
- **Codeine and Oxycodone** should be avoided. They can cause excessive sleepiness in babies.
- The FDA has advised breastfeeding mothers to NOT take **Tramadol**.
- **Ibuprofen or Tylenol**, taken as directed, are safe choices for pain control while breastfeeding.

You or your physician may access the latest information available regarding any specific drug at:

<https://toxnet.nlm.nih.gov/newtoxnet/lactmed.htm> or <https://www.ncbi.nlm.nih.gov/books/NBK501922/>

Reference used: ABM Clinical Protocol #15: Analgesia and Anesthesia for the Breastfeeding Mother, Revised 2017. <https://abm.memberclicks.net/assets/DOCUMENTS/PROTOCOLS/15-analgesia-and-anesthesia-protocol-english.pdf>

Reviewed: copyright March 2020

MilkWorks



Community Breastfeeding Center

5930 S. 58th Street
(in the Trade Center)
Lincoln, NE 68516
(402) 423-6402

10818 Elm Street
Rockbrook Village
Omaha, NE 68144
(402) 502-0617

For additional
information:
www.milkworks.org